

University of Groningen

Molecular heterogeneity and angiogenesis in glioblastoma

Conroy, Siobhan

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2017

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Conroy, S. (2017). *Molecular heterogeneity and angiogenesis in glioblastoma*. [Thesis fully internal (DIV), University of Groningen]. Rijksuniversiteit Groningen.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Chapter 13

Summary, discussion and future perspectives



Summary and discussion

Glioblastoma (GBM), a highly malignant brain neoplasm, represents one of the most complex types of cancer. The diagnosis GBM leaves patients with a dismal prognosis and limited therapeutic options are available, which all can only transiently halt disease progression. Better understanding of the genetic machinery that results in this devastating disease is hoped to improve these perspectives and over the last decade the molecular classification of GBMs has moved on apace. During the initiation of the studies described in the thesis, molecular subclasses based on transcriptional profiles had just been identified [1-3], and a vast majority of studies evaluating the druggable targets within these subclasses was underway.

With **Part I** of the thesis we have aimed to contribute to this understanding by performing several studies on GBM tumor material that was surgically resected from patients. First, we explored the possibility of molecular subclassification of GBMs in **Chapter 3** on tumor material that is already routinely obtained for diagnosing patients, and with the use of only widely used pathological methods. To prevent using up the available tissue and to allow for the analyses of an extensive panel of markers, we created a tissue microarray that contained GBM tumor tissue from 167 patients. Through the application of primarily immunohistochemical stainings, we were able to subclassify two-thirds of the tumors into proneural (PN), classical (CLAS) and mesenchymal (MES) subclasses.

In **Chapter 5** and **Chapter 6** we assessed the influence of environmental stimuli on subclass identity of GBM tumor cells. Glioma stem-like cells (GSCs) are endowed with tumor-initiating properties [4] and some studies have shown the role of GSCs in clinical relapse due to therapy refractoriness [5-7]. Microenvironmental support is provided to the GSCs as they reside in their preferential perivascular niche [8], and we showed that both TGF- β and hypoxia can induce the expression of MES markers in glioma cell models. This should be interpreted as malignant progression, since a PN-to-MES transition has been observed at disease recurrence and the MES phenotype has been reported to mediate radio-resistance [2,7]. These studies indicate that environmental cues, as we also discussed extensively in **Chapter 2**, can contribute significantly to GBM progression.

Due to extensive associations between receptors with tyrosine kinase domains and the above-described molecular subclasses, we explored kinase activity profiles of molecular subclasses in **Chapter 4**. Unexpectedly, only few phosphorylation sites on the array were found to associate with molecular subclasses, irrespective of the mutational status of, for example, GBMs of the CLAS subtype. This suggests that although receptor expression levels are aberrant in subgroups of tumors, substantial ongoing cross-talk between multiple signaling routes in GBMs prevented the identification of canonically exploited signaling pathways. It is also possible that the substantial percentage of non-tumor cells in GBM tissue lysates desensitized the array, and together these findings complicate therapeutic target selection in GBM further.

Part II of the thesis then shifted focus to the role of angiogenesis in GBM. The examination

of vascular parameters in **Chapter 8** on GBM tissue sections indicated that the ascribed angiogenic profile to the MES subclass could have been confounded by the fact that these tumors had a similar number of vessels, but that these were merely larger in size. It should be noted that the functional interpretation would benefit greatly from insight into the perfusion status, because pro-thrombotic mechanisms [9], vascular normalization [10,11] and edema [10] represent significant modifiers of vascular effects.

In **Chapter 9** we subsequently identified IL-8 as a potentially important mediator of the vascular enlargement that we observed in MES GBM tissue from patients. IL-8 was substantially higher expressed in pro-angiogenic tumor, MES GSC lines and the addition of an IL-8 neutralizing antibody only reduced angiogenesis when it was added to the conditioned medium of MES GSCs. It could however be noted that the *in vitro* results in this chapter are limited in effect size, but the differences in, for example, *in vitro* serum requirements between cell lines severely impeded absolute quantitation of paracrine effects and could have prevented the identification of actual larger effect sizes. The exploration of IL-8 inhibition in an *in vivo* xenograft model comparing the effect of treatment on engrafted PN and MES GSCs could substantiate our findings further.

In contrast to the microenvironmental approach adopted in the studies where subclass plasticity was assessed, we next examined the effects of key angiogenic mediators with a strong focus on tumor cell altering effects instead. Angiopoietin-2 (Ang-2) and VEGFA, well-known for their pro-angiogenic effects in glioma angiogenesis [12], exerted an unexpected indirect anti-angiogenic effect when they were used to stimulate tumor cells *in vitro* as is described in **Chapter 10**. *In vivo* stimulation with these agents, however, did not alter tumor growth or vascularization parameters, potentially signifying the inability of angiogenesis modulation in xenograft GBM models. In contrast, the inhibition of Ang-2 and VEGFA in **Chapter 11** in the exact same model did result in reduced expression of vascular parameters.

The discrepancies between these studies could potentially be explained by several factors. First, the fact that an anti-angiogenic effect with conditioned medium from GBM cells is observed relative to the control PBS-stimulation condition, perhaps needs to be rephrased to reduced pro-angiogenic effects of GBM cells on endothelium. Second, the expression analyses of angiogenic signaling molecules in **Chapter 8** and **Chapter 9** illustrate that a multitude of signaling routes are exploited in GBMs, which allows the xenografts to already reach a vascular saturation stage even before application of our treatments. It is possible that the selection of this xenograft GBM model prevented the detection of the anti-angiogenic properties of Klotho in **Chapter 12**.

Two recent reports allow us to compare the combined Ang-2 and VEGFA inhibition that we performed in our immunocompromised model to studies that were performed in immunocompetent GBM models. The reports of the other studies indicated that the recruitment of macrophages and the phenotypical transition of these following combined Ang-2 and VEGFA inhibition served a crucial role in the survival extension that was achieved in their *in vivo* studies [13,14]. Since we were unable to identify an effect on tumor growth or survival, we conclude that the immunocompromised environment under which our *in vivo* experiment was performed likely contributed substantially to treatment outcome. The multifaceted properties of both Ang-2 and VEGFA are also extensively reviewed in **Chapter 7**, and we speculate that

the pro-inflammatory effects of especially Ang-2 likely have more potential in an anti-tumor approach than the combined anti-angiogenesis approach that was employed by us [13-15].

Future perspectives

Although we want to acknowledge the substantial advancements in our understanding of the elusive complexity and molecular heterogeneity of GBMs by The Cancer Genome Atlas initiative and associated groups, the therapeutic implications of the extensive molecular classification have thus far not been groundbreaking. Apart from selective benefit from the anti-angiogenic drug Bevacizumab in PN tumors that do not harbor a mutation in the *isocitrate dehydrogenase 1* gene (*IDH1*, [16]), to our knowledge no selective therapeutic benefit of molecular GBM subclasses has been reported. An ongoing clinical trial that specifically focuses on *epidermal growth factor receptor* (*EGFR*) amplification, an aberration associated with the transcriptionally described CLAS subgroup, is another example of a trial that is at an advanced clinical stage, but notably also addresses a specific molecular aberration and not an expression pattern. Clinically a tendency can thus be observed that specific aberrations are targeted instead of an expression signature. Additionally, the field has progressed with the identification of molecular heterogeneity of GBMs in methylation patterns [17] and genomic profiles that independently resulted in an alternative GBM classification [18]. The aberrations described in the latter study have recently been adopted by the new WHO guidelines [19] and studies to be initiated will very likely focus on these aberrations.

By focusing on specific molecular aberrations as described in the 2016 WHO guidelines, it is ensured that tumor specific alterations are studied, given that molecular aberrations in these tumors generally do not concern somatic mutations. The increase in GBM incidence with age also argues for that notion [20]. The morphometrical analyses of the tumor vasculature in this thesis and the high percentage of myeloid cell content in freshly isolated biopsies both call into question whether transcriptional profiling of whole GBM tissue that exhibits such cellular heterogeneity is appropriate and in many ways has allowed us to appreciate another dimension of multiform in GBM. Lower tumor cell percentages could have masked many small but important tumor specific properties.

The discordant results obtained with *in vivo* experimentation in this thesis shed light on the important differences between *in vitro*, *in ovo* and *in vivo* models that were employed for our experiments. It underscores the artificial aspect of *in vitro* culturing, as we are, for example, simply unable to define whether conditioned medium from GBM tumor cells exerted a pro-angiogenic effect due to the extensive set of experimental parameters that are controlled. Unfortunately this controlled experimental environment is not inspired by physiological or clinical considerations, but merely to allow for the observation of consistent effects. Improved *in vitro* models that would allow more physiological (co-)culturing conditions could help us gain insight into the crucial interactions that ultimately contribute to tumor progression. Complementary approaches involving both clinically relevant xenograft models alongside

immunocompetent transgenic mouse models will be needed to reach definitive conclusions on the role the microenvironment in GBM.

As a concluding remark, I believe that the field could advance by investing more in the study of therapy refractoriness. The distressing situation that patients with GBM undergo fully justifies the exploration of novel, currently unapproved therapies for this disease. From a research perspective, however, a balance is required between the exploration of novel promising agents and the attempts to understand why other agents did not provide benefit, because the latter could very well be our next step to improved therapeutic efficacy. To accomplish this, advances in understanding of basic tumor biology, plasticity, therapy response and resistance are fundamental.

References

1. Freije WA, Castro-Vargas FE, Fang Z, Horvath S, Cloughesy T, Liao LM, Mischel PS, Nelson SF. Gene expression profiling of gliomas strongly predicts survival. *Cancer Res.* 2004;64(18):6503-6510.
2. Phillips HS, Kharbanda S, Chen R, Forrest WF, Soriano RH, Wu TD, Misra A, Nigro JM, Colman H, Soroceanu L, Williams PM, Modrusan Z, Feuerstein BG, Aldape K. Molecular subclasses of high-grade glioma predict prognosis, delineate a pattern of disease progression, and resemble stages in neurogenesis. *Cancer Cell.* 2006;9(3):157-173.
3. Verhaak RG, Hoadley KA, Purdom E, Wang V, Qi Y, Wilkerson MD, Miller CR, Ding L, Golub T, Mesirov JP, Alexe G, Lawrence M, O'Kelly M, Tamayo P, Weir BA, Gabriel S, Winckler W, Gupta S, Jakkula L, Feiler HS, Hodgson JG, James CD, Sarkaria JN, Brennan C, Kahn A, Spellman PT, Wilson RK, Speed TP, Gray JW, Meyerson M, Getz G, Perou CM, Hayes DN, Cancer Genome Atlas Research Network. Integrated genomic analysis identifies clinically relevant subtypes of glioblastoma characterized by abnormalities in PDGFRA, IDH1, EGFR, and NF1. *Cancer Cell.* 2010;17(1):98-110.
4. Singh SK, Hawkins C, Clarke ID, Squire JA, Bayani J, Hide T, Henkelman RM, Cusimano MD, Dirks PB. Identification of human brain tumour initiating cells. *Nature.* 2004;432(7015):396-401.
5. Eramo A, Ricci-Vitiani L, Zeuner A, Pallini R, Lotti F, Sette G, Piloizzi E, Larocca LM, Peschle C, De Maria R. Chemotherapy resistance of glioblastoma stem cells. *Cell Death Differ.* 2006;13(7):1238-1241.
6. Bao S, Wu Q, McLendon RE, Hao Y, Shi Q, Hjelmeland AB, Dewhirst MW, Bigner DD, Rich JN. Glioma stem cells promote radioresistance by preferential activation of the DNA damage response. *Nature.* 2006;444(7120):756-760.
7. Bhat KP, Balasubramanian V, Vaillant B, Ezhilarasan R, Hummelink K, Hollingsworth F, Wani K, Heathcock L, James JD, Goodman LD, Conroy S, Long L, Lelic N, Wang S, Gumin J, Raj D, Kodama Y, Raghunathan A, Olar A, Joshi K, Pelloso CE, Heimberger A, Kim SH, Cahill DP, Rao G, Den Dunnen WF, Boddeke HW, Phillips HS, Nakano I, Lang FF, Colman H, Sulman EP, Aldape K. Mesenchymal differentiation mediated by NF-kappaB promotes radiation resistance in glioblastoma. *Cancer Cell.* 2013;24(3):331-346.
8. Calabrese C, Poppleton H, Kocak M, Hogg TL, Fuller C, Hamner B, Oh EY, Gaber MW, Finklestein D, Allen M, Frank A, Bayazitov IT, Zakharenko SS, Gajjar A, Davidoff A, Gilbertson RJ. A perivascular niche for brain tumor stem cells. *Cancer Cell.* 2007;11(1):69-82.
9. Brat DJ, Van Meir EG. Vaso-occlusive and prothrombotic mechanisms associated with tumor hypoxia, necrosis, and accelerated growth in glioblastoma. *Lab Invest.* 2004;84(4):397-405.

10. Batchelor TT, Sorensen AG, di Tomaso E, Zhang WT, Duda DG, Cohen KS, Kozak KR, Cahill DP, Chen PJ, Zhu M, Ancukiewicz M, Mrugala MM, Plotkin S, Drappatz J, Louis DN, Ivy P, Scadden DT, Benner T, Loeffler JS, Wen PY, Jain RK. AZD2171, a pan-VEGF receptor tyrosine kinase inhibitor, normalizes tumor vasculature and alleviates edema in glioblastoma patients. *Cancer Cell*. 2007;11(1):83-95.
11. Batchelor TT, Gerstner ER, Emblem KE, Duda DG, Kalpathy-Cramer J, Snuderl M, Ancukiewicz M, Polaskova P, Pinho MC, Jennings D, Plotkin SR, Chi AS, Eichler AF, Dietrich J, Hochberg FH, Lu-Emerson C, Iafrate AJ, Ivy SP, Rosen BR, Loeffler JS, Wen PY, Sorensen AG, Jain RK. Improved tumor oxygenation and survival in glioblastoma patients who show increased blood perfusion after cediranib and chemoradiation. *Proc Natl Acad Sci U S A*. 2013;110(47):19059-19064.
12. Hardee ME, Zagzag D. Mechanisms of glioma-associated neovascularization. *Am J Pathol*. 2012;181(4):1126-1141.
13. Peterson TE, Kirkpatrick ND, Huang Y, Farrar CT, Marijt KA, Kloepper J, Datta M, Amoozgar Z, Seano G, Jung K, Kamoun WS, Vardam T, Snuderl M, Goveia J, Chatterjee S, Batista A, Muzikansky A, Leow CC, Xu L, Batchelor TT, Duda DG, Fukumura D, Jain RK. Dual inhibition of ang-2 and VEGF receptors normalizes tumor vasculature and prolongs survival in glioblastoma by altering macrophages. *Proc Natl Acad Sci U S A*. 2016;113(16):4470-4475.
14. Kloepper J, Riedemann L, Amoozgar Z, Seano G, Susek K, Yu V, Dalvie N, Amelung RL, Datta M, Song JW, Askoxylakis V, Taylor JW, Lu-Emerson C, Batista A, Kirkpatrick ND, Jung K, Snuderl M, Muzikansky A, Stubenrauch KG, Krieter O, Wakimoto H, Xu L, Munn LL, Duda DG, Fukumura D, Batchelor TT, Jain RK. Ang-2/VEGF bispecific antibody reprograms macrophages and resident microglia to anti-tumor phenotype and prolongs glioblastoma survival. *Proc Natl Acad Sci U S A*. 2016;113(16):4476-4481.
15. Scholz A, Plate KH, Reiss Y. Angiopoietin-2: A multifaceted cytokine that functions in both angiogenesis and inflammation. *Ann N Y Acad Sci*. 2015;1347:45-51.
16. Sandmann T, Bourgon R, Garcia J, Li C, Cloughesy T, Chinot OL, Wick W, Nishikawa R, Mason W, Henriksson R, Saran F, Lai A, Moore N, Kharbanda S, Peale F, Hegde P, Abrey LE, Phillips HS, Bais C. Patients with proneural glioblastoma may derive overall survival benefit from the addition of bevacizumab to first-line radiotherapy and temozolomide: Retrospective analysis of the AVAglio trial. *J Clin Oncol*. 2015;33(25):2735-2744.
17. Noushmehr H, Weisenberger DJ, Diefes K, Phillips HS, Pujara K, Berman BP, Pan F, Pelloso CE, Sulman EP, Bhat KP, Verhaak RG, Hoadley KA, Hayes DN, Perou CM, Schmidt HK, Ding L, Wilson RK, Van Den Berg D, Shen H, Bengtsson H, Neuvial P, Cope LM, Buckley J, Herman JG, Baylin SB, Laird PW, Aldape K, Cancer Genome Atlas Research Network. Identification of a CpG island methylator phenotype that defines a distinct subgroup of glioma. *Cancer Cell*. 2010;17(5):510-522.
18. Eckel-Passow JE, Lachance DH, Molinaro AM, Walsh KM, Decker PA, Sicotte H, Pekmezci M, Rice T, Kosel ML, Smirnov IV, Sarkar G, Caron AA, Kollmeyer TM, Praska CE, Chada AR, Halder C, Hansen HM, McCoy LS, Bracci PM, Marshall R, Zheng S,

- Reis GF, Pico AR, O'Neill BP, Buckner JC, Giannini C, Huse JT, Perry A, Tihan T, Berger MS, Chang SM, Prados MD, Wiemels J, Wiencke JK, Wrensch MR, Jenkins RB. Glioma groups based on 1p/19q, IDH, and TERT promoter mutations in tumors. *N Engl J Med*. 2015;372(26):2499-2508.
19. Louis DN, Perry A, Reifenberger G, von Deimling A, Figarella-Branger D, Cavenee WK, Ohgaki H, Wiestler OD, Kleihues P, Ellison DW. The 2016 world health organization classification of tumors of the central nervous system: A summary. *Acta Neuropathol*. 2016;131(6):803-820.
 20. Ostrom QT, Gittleman H, Xu J, Kromer C, Wolinsky Y, Kruchko C, Barnholtz-Sloan JS. CBTRUS statistical report: Primary brain and other central nervous system tumors diagnosed in the united states in 2009-2013. *Neuro Oncol*. 2016;18:v1-v75.

